

TB SKIN TEST/SCREENING DOCUMENTATION FORM

Check applicable:

- Initial two-step TST (Mantoux) for all staff having direct contact with patients/ clients
 - Step 1- Initial TST
 - Step 2- TST 1-3 weeks after step 1
- OR
- Documentation of negative TST within 12 months prior to hire, and
 - One-step TST
- Annual (all employees providing patient/ client care)
 - Symptom Screen
- Biannual (for health care workers frequently exposed)
 - Symptom Screen
- Post-exposure
 - Administer TST as soon as possible after exposure,
 - If initial post exposure TST is negative, repeat at 8-10 weeks post-exposure
- Previously documented positive TST
- Previously infected with nontuberculosis mycobacterium
- Radiograph excluding TB Disease

SKIN TEST

Step #1 _____ was given a Mantoux tuberculin ppd intradermal skin test by _____ on _____ on left/right forearm.

Lot# _____ Brand _____ Expiration _____
Results _____ mm induration Date _____ Read by _____

Step #2 by _____ on _____ on left/right forearm.

Lot# _____ Brand _____ Expiration _____
Results _____ mm induration Date _____ Read by _____

Note: Do not include redness or ulceration reading your results. Read results across (transverse) forearm. See TB protocol for classification of results.

If a skin test is > 10mm and has one or more risk factors for infection, the employee should be referred to the county health department or a local physician for follow-up assessment.

Employee Name: _____

Signature: _____ Date: _____